

THE ANTHROPOCENE by the numbers

Griffin Chure¹, Rachel A. Banks², Avi Flamholz², Nicholas S. Sarai³, Mason Kamb⁴, Ignacio Lopez-Gomez⁵, Yinon Bar-On⁶, Ron Milo⁶, Rob Phillips^{2,7,*}

California Institute of Technology, Pasadena, CA, USA, 91125:

1. Department of Applied Physics; 2. Division of Biology and Biological Engineering; 3. Division of Chemistry and Chemical Engineering
 4. Chan-Zuckerberg BioHub, San Francisco, CA, 94158
 5. Department of Environmental Science and Engineering; 7. Department of Physics;
 6. Weizmann Institute of Science, Rehovot 7610001, Israel; Department of Plant and Environmental Sciences

ABSTRACT

A candidate for the greatest experiment of the last 10,000 years is the presence and action of modern human beings on planet Earth, the often complex results of which are now being felt on various fronts. While there has been a deluge of careful studies exploring each facet of these "human impacts" on Earth, they are often highly focused and necessarily technical, rarely displaying their integration with other human impacts as a whole. In this snapshot, we present a diverse (yet necessarily incomplete) array of quantities that summarize the broad reach of human action across the planet.

CATEGORIES
LAND
ENERGY
WATER
FLORA & FAUNA
ATMOSPHERIC & BIOGEOCHEMICAL CYCLES

REFERENCE UNITS		
human population $\approx 8 \times 10^9$		
area		
soccer field	tennis court	room
$\approx 7000 \text{ m}^2$	$\approx 300 \text{ m}^2$	$\approx 10 \text{ m}^2$
volume		
shipping container	bathub	incandescent lightbulb
$\approx 30 \text{ m}^3$	$\approx 0.1 \text{ m}^3$	$\approx 100 \text{ W}$
mass		
pick-up truck	refrigerator	cinder block
$\approx 2000 \text{ kg}$	$\approx 100 \text{ kg}$	$\approx 10 \text{ kg}$

C SEA ICE EXTENT
 sea ice extent loss $\approx 5 \times 10^{10} \text{ m}^2 \approx \frac{1}{2}$ per per yr

D MATERIAL PRODUCTION
 concrete $\approx 3 \times 10^{13} \text{ kg} / \text{yr} \approx 2$ per per yr
 steel $\approx 2 \times 10^{12} \text{ kg} / \text{yr} \approx 2.5$ per per yr
 plastic $\approx 4 \times 10^{11} \text{ kg} / \text{yr} \approx 5$ per per yr

E LIVESTOCK POPULATION
 chicken $\approx 2.5 \times 10^{10} \approx 3$ per per yr
 cattle $\approx 1.5 \times 10^9 \approx 1$ per 5
 swine $\approx 1 \times 10^9 \approx 1$ per 8
 all livestock $\approx 3 \times 10^{10} \approx 4$ per per yr

F NITROGEN FIXATION
 synthetic nitrogen fixation $\approx 1.5 \times 10^{11} \text{ kg} \approx 2$ per per yr

G OCEAN ACIDITY
 % increase in H^+ concentration $\approx 0.4\%$ per yr

H HUMAN POPULATION
 urban $\approx 4.5 \times 10^9$
 rural $\approx 3.5 \times 10^9$
 total $\approx 8 \times 10^9$

I GREENHOUSE GAS EMISSIONS
 $\text{CO}_2 \approx 4 \times 10^{13} \text{ kg} \approx 2.5$ per per yr
 $\text{CH}_4 \approx 4 \times 10^{11} \text{ kg} \approx 5$ per per yr
 $\text{N}_2\text{O} \approx 1 \times 10^{10} \text{ kg} \approx 1$ per 10 per yr

J SEA LEVEL RISE
 rise due to added water $\approx 2 \text{ mm}$ per yr
 rise due to thermal expansion $\approx 1.2 \text{ mm}$ per yr
 total rise $\approx 3.3 \text{ mm}$ per yr

K OCEAN WARMING
 heat uptake $\approx 350 \text{ TW} \approx 435$ per
 temperature change in top 700 m since 1960 $\approx 0.2 \text{ }^\circ\text{C}$

L NUCLEAR FALLOUT
 globally-distributed mass of plutonium from $\approx 3300 \text{ kg} \approx 1.5$ per per yr

M CONTEMPORARY EXTINCTION
 animal species extinct since 1500 > 750
 plant species extinct since 1500 > 120

N EARTH MOVING
 waste from coal mining $\approx 7 \times 10^{13} \text{ kg} / \text{yr} \approx 46$ per per yr
 urbanization $> 15 \times 10^{13} \text{ kg} / \text{yr} \approx 9$ per per yr
 agriculture $\approx 2 \times 10^{13} \text{ kg} / \text{yr} \approx 1$ per per yr

G LAND USE
 agriculture $\approx 5 \times 10^{13} \text{ m}^2 \approx 1$ per per yr
 urban $\approx 7 \times 10^{11} \text{ m}^2 \approx \frac{1}{5}$ per per yr

H WATER WITHDRAWAL
 agriculture $\approx 1.3 \times 10^{12} \text{ m}^3 / \text{yr} \approx 6$ per per yr
 industrial $\approx 6 \times 10^{11} \text{ m}^3 / \text{yr} \approx 2.5$ per per yr
 domestic $\approx 5 \times 10^{10} \text{ m}^3 / \text{yr} \approx 65$ per per yr
 total $\approx 2 \times 10^{12} \text{ m}^3 / \text{yr} \approx 8$ per per yr

I TOTAL POWER USE
 global $\approx 18 \text{ TW} \approx 23$ per

J POWER FROM FOSSIL FUELS
 natural gas $\approx 4.5 \text{ TW} \approx 6$ per
 coal $\approx 5 \text{ TW} \approx 7$ per
 oil $\approx 6.5 \text{ TW} \approx 8$ per
 total $\approx 16 \text{ TW} \approx 20$ per


K POWER FROM RENEWABLES
 wind $\approx 0.3 \text{ TW} \approx 1$ per 3
 solar $\approx 0.2 \text{ TW} \approx 1$ per 4
 hydroelectric $\approx 1 \text{ TW} \approx 1.5$ per
 total $\approx 2 \text{ TW} \approx 2.5$ per

L POWER FROM NUCLEAR FISSION
 nuclear power $\approx 0.8 \text{ TW} \approx 1$ per

M RIVER FRAGMENTATION
 volume of dammed and fragmented rivers $\approx 6 \times 10^{11} \text{ m}^3 \approx 3$ per per yr

N TREE COVERAGE AREA LOSS
 deforestation $\approx 6 \times 10^{10} \text{ m}^2 \approx \frac{3}{4}$ per per yr
 forestry $\approx 5 \times 10^{10} \text{ m}^2 \approx \frac{1}{2}$ per per yr
 wildfire $\approx 7 \times 10^{10} \text{ m}^2 \approx 1$ per per yr
 agriculture $\approx 7 \times 10^{10} \text{ m}^2 \approx 1$ per per yr
 urbanization $\approx 2 \times 10^9 \text{ m}^2 \approx 1$ per 40 per yr
 total $\approx 2 \times 10^{11} \text{ m}^2 \approx 2.5$ per per yr

O FOSSIL FUEL EXTRACTION
 natural gas $\approx 4 \times 10^{12} \text{ m}^3 / \text{yr} \approx 16$ per per yr
 oil $\approx 6 \times 10^9 \text{ m}^3 / \text{yr} \approx 8$ per per yr
 coal $\approx 8 \times 10^{12} \text{ kg} / \text{yr} \approx \frac{1}{2}$ per per yr




oil & natural gas mineral resources
 coal mineral resources

The Anthropocene by the Numbers — Impacts By Region


THE GEOGRAPHY OF HUMAN IMPACTS

Page 1 represents the impact humans have on the Earth at a global scale. While these numbers are handy, it is important to acknowledge that they vary from country-to-country and continent-to-continent. Furthermore, the consequences of these anthropogenic impacts are also unequally distributed, meaning some regions experience effects disproportionate to their contribution. Here, we give a sense of the geographic distribution of several values presented on page 1, broken down by continental region as shown below.



THE LIVESTOCK POPULATION


The global population of terrestrial livestock is around 30 billion individuals, most of which are chickens. Asia houses most of the global livestock population, though South America and Europe harbor more animals on a per-capita basis.



Sources: Food and Agricultural Organization of the United Nations

NITROGENOUS FERTILIZER USE & PRODUCTION


Modern agriculture requires nitrogen in amounts beyond what is produced naturally. Asia synthesizes and consumes a large majority of fixed nitrogen. However, Europe and North America dominate per capita synthesis whereas Oceania consumes more fertilizer per capita than any other region.



Sources: Food and Agricultural Organization (FAO) of the United Nations. Notes: Values account for reactive nitrogen production/consumption in context of fertilizer only and does not account for plastics, explosives, or other uses.

THE HUMAN POPULATION


There are ≈ 8 billion humans on the planet, with approximately 50% living in 'urban' environments. The majority of the world's population (as well as the majority of both urban and rural dwellers) live in Asia.



Sources: Food and Agricultural Organization of the United Nations - World Population Notes: Urban/rural designation has no set definition and follows the conventions set by each reporting country.

WATER WITHDRAWAL


While Asia withdraws the most water for agricultural and municipal needs, North America withdraws the plurality of water for industrial purposes. North America also withdraws more water per capita than any other region.



Sources: AQUASTAT Main Database. Food and Agriculture Organization of the United Nations. Notes: Values are reported directly from member countries and represent average of 2013-2017 period. Per capita values are computed given population of reporting countries.

GREENHOUSE GAS EMISSIONS

CO₂ and CH₄ are two potent greenhouse gases which are routinely emitted by anthropogenic processes such as burning fuel and rearing livestock. While Asia emits roughly half of all CO₂ and CH₄, North America and Oceania produce the most on a per capita basis, respectively.




Sources: CO₂ data collated by: Friedlingstein, P. et al. (2019). doi: 10.5194/essd-11-1783-2019. See Panel J on Pg. 4 for complete list of sources. CH₄ data from Saunios et al, 2020 doi: 10.5194/essd-12-1561-2020 Notes: Values report decadal averages in kg CO₂ or CH₄ per year over time period 2008-2017.

POWER GENERATION AND CONSUMPTION


From heating water, to powering lights, to moving our vehicles, nearly every facet of modern human life requires the consumption of power, culminating in nearly 20 TW of power use in recent years. Asia consumes over half of the power derived from combustion of fossil fuels, with Europe and North America each consuming around 20% of the global total. Asia also produces the plurality of power from renewable technologies, such as hydroelectric, wind, and solar, however, North America, South America, and Europe each produce more on a per capita basis. Nuclear energy, however, is primarily produced in Europe, with North America and Asia coming in second and third place, respectively. On a per-capita basis, North America consumes or produces more energy than all other regions considered here, yielding a total power consumption of nearly 10,000 W per person.

Sources: Energy Information Administration of the United States (2017) Notes: "Renewables" includes hydroelectric, biofuels, biomass (wood), geothermal, wind, and solar. "Fossil fuels" includes coal, oil, and natural gas.



LAND USE

Though humans are nearly evenly split between urban and rural environments, agricultural land is the far more common use of land area. Together, Asia and Africa contain more than half of global agricultural land. Asia alone accommodates more than half of the global urban land area.




Sources: Food and Agricultural Organization (FAO) of the United Nations (2015) - Land Use [agricultural area], Florczyk et al. 2019 - GHS Urban Centre Database 2015 [urban land area]. Notes: Urban is defined as any inhabited area with ≥ 2500 residents, as defined by the USDA.

TREE COVERAGE AREA LOSS

Most drivers of tree coverage area loss are comparable in their effect at a global scale. However, there are drastic regional differences in the relative magnitudes.

REGION DEFINITION


Central & South America, Russia, China, & South Asia
 North America, Southeast Asia, Europe (- Russia)
 Africa, Oceania



Source: Curtis et al. 2018 doi: 10.1126/science.aau3445. Notes: Regions are as reported in Curtis et al. 2018. "Deforestation" here denotes permanent removal of tree cover for commodity production. "Shifting agriculture" here denotes forest/shrub land converted to agriculture and later abandoned. All values correspond to breakdown of cumulative tree cover area loss from 2001 - 2015.

MATERIAL PRODUCTION

Humans excavate an enormous amount of material from the Earth's crust and transform it to build our structures. Two of these materials, concrete and steel, are produced primarily in Asia on both a global and per capita basis. Asia's per capita production of steel is only outpaced by Europe.




Sources: USGS Statistics and Information 2020, Steel Statistical Yearbook 2019 World Steel Association. Food and Agricultural Organization (FAO) of the United Nations - Annual Population. Notes: Reported values for concrete and steel production corresponds to 2017 and 2018 values, respectively. Mass of concrete was calculated using a rule-of-thumb that 1 kg of cement yields 7 kg of concrete (Monteiro et al. 2017. doi: 10.138/nmat4930).

The Anthropocene by the Numbers — Dimensionless Ratios

Much of our understanding of the scales of things is comparative. When we measure lengths, we do so relative to some intuitive distance that provides context. Our aim here is to present some “yardstick” to measure those numbers presented on pages 1 and 2 in a ratiometric form that compares the magnitude of a given human impact to a natural scale for that same quantity. For example, in considering the use of land by humans, a natural dimensionless way to characterize that number is by comparing it to the total land area of our planet, a comparison that yields what we call the “Terra number.”


Similarly, when we consider the entirety of human-made materials, it is natural to compare this mass to the total biomass on our planet. Here we present twelve key human impacts in this dimensionless form. These numbers describe the solid earth, the atmosphere, the biosphere, the oceans, and human resource and energy use, and we hope that our readers will be emboldened to consider their own favorite examples in a similar dimensionless format. Where appropriate, we reference key values using a [Human Impacts Database](#) number (HuID) accessible via [anthroponumbers.org](#)

THE TERRA NUMBER

$$Te = \frac{\text{land area used by humans}}{\text{total land area of Earth}} \approx 0.3$$



The Terra Number captures the extent to which we have taken control of Earth’s terrestrial surface to support our dwellings and, more importantly, our agriculture. Of the $\approx 1.5 \times 10^{14} \text{ m}^2$ of Earth’s surface area that is land, approximately $5 \times 10^{13} \text{ m}^2$ (HuID: 29582) is used for agriculture, including growing our crops and rearing livestock. Despite being icons of humanity, urban centers occupy between 6.5 and $7.5 \times 10^{13} \text{ m}^2$ (HuID: 41339, 39341), a total less than 1% of the terrestrial surface. In total, humans directly manage 30% of Earth’s terrestrial surface.

THE DEFORESTATION NUMBER

$$Df = \frac{\text{annual forest loss from human action}}{\text{annual forest loss from wildfire}} \approx 2$$



The Deforestation Number reflects the magnitude to which we intentionally clear land of tree cover for production of goods, agriculture, and building our cities relative to the tree cover lost due to wildfire. The intentional clearing of land for production of goods (such as lumber and paper) permanently deforests $\approx 6 \times 10^{10} \text{ m}^2$ per year. Other intentionally cleared area (which eventually regrows) is comparable in magnitude with $\approx 5 \times 10^{10} \text{ m}^2 / \text{yr}$ for forestry (HuID: 38352) and $\approx 7.5 \times 10^{10} \text{ m}^2 / \text{yr}$ for shifting agriculture (HuID: 24388). Expansion of urban areas accounts for $< 1\%$ of the total annual deforested area, averaging $\approx 2 \times 10^9 \text{ m}^2 / \text{yr}$ (HuID: 19429). In total, intentional deforestation by humans amounts to $\approx 1.5 \times 10^{11} \text{ m}^2 / \text{yr}$, about twice the area cleared by natural and human-caused wildfires each year ($\approx 7 \times 10^{10} \text{ m}^2 / \text{yr}$, HuID: 92221).

THE BARNYARD NUMBER

$$By = \frac{\text{mass of terrestrial livestock}}{\text{mass of terrestrial wild animals}} \approx 30$$



The Barnyard Number focuses another lens onto the massive agricultural transformation of the planet by comparing the total biomass of terrestrial livestock (e.g. cows, chickens, and pigs) to that of terrestrial wild mammals and birds (e.g. elephants, foxes, and pelicans) [1]. Agricultural intensification of the 20th century has resulted in livestock outweighing all wild terrestrial animals by a factor of ≈ 30 . While poultry make up the vast majority of terrestrial livestock (≈ 25 billion individuals, HuID: 94934), they represent a small proportion of livestock biomass. Despite a smaller population of ≈ 1.5 billion (HuID: 92006), cattle dominate livestock biomass with a total mass of $\approx 1.5 \times 10^{12} \text{ kg}$.

THE NITROGEN NUMBER

$$N_2 = \frac{\text{mass of } N_2 \text{ fixed via the Haber-Bosch Process}}{\text{mass of } N_2 \text{ fixed biologically}} \approx 1$$



The Nitrogen Number illustrates how humans have transformed the global nitrogen cycle to sustain a global population in excess of three billion humans. While molecular nitrogen (N_2) is abundant in our atmosphere, nitrogen can only be used by plants in a reactive form such as ammonia (NH_3). The 1910 development of the Haber-Bosch process for industrial synthesis of NH_3 from N_2 was critical for supporting the agricultural needs of a growing human population and for supplying NH_3 for chemical and explosive synthesis. Primarily through the Haber-Bosch process, humans synthesize as much reactive nitrogen industrially ($\approx 1.5 \times 10^{11} \text{ kg / yr}$, HuID: 61614, 60580) as is synthesized by nitrogen-fixing microbes in terrestrial ecosystems ($\approx 1 \times 10^{11} \text{ kg per year}$, HuID: 15205). Beyond influencing the environmental balance of reactive nitrogen, modern synthesis technologies require a sizable amount of energy, contributing significantly to global CO_2 emissions.

THE WATER NUMBER

$$Wa = \frac{\text{annual water volume used by humans}}{\text{global annual river discharge volume}} \approx 0.05$$



The Water Number captures the magnitude of human water usage relative to global river discharge, a major source of renewable freshwater. Agriculture defines this aspect of human impacts, using $\approx 1.5 \times 10^{12} \text{ m}^3$ (HuID: 43593) of water annually, accounting for the majority of human water usage. Water used for industrial purposes, including cooling thermoelectric plants amounts to $5.9 \times 10^{11} \text{ m}^3 / \text{yr}$ (HuID: 27142), and domestic use is $\approx 6 \times 10^{10} \text{ m}^3 / \text{yr}$ (HuID: 69424). In total, annual human water withdrawal is about 5% of global annual river discharge volume, a major source of renewable freshwater. While this is a small fraction, available freshwater is highly variable across the globe and about a third of the human population lives in water stressed areas, where greater than 40% of available freshwater is used.

THE RIVER NUMBER

$$Rv = \frac{\text{river volume controlled by humans}}{\text{free-flowing river volume}} \approx 1$$



The River Number illustrates the extent to which we have fragmented the free-flowing river systems of the globe for irrigation, flood control, and generation of hydroelectric power. Harnessing this water, however, requires damming the river – thus interrupting its flow and altering the riverine ecosystem. Primarily through damming and construction of channels, humans control $\approx 6 \times 10^{11} \text{ m}^3$ of water (HuID: 61661), a volume comparable to that freely flowing in unperturbed river systems [2]. Of the free-flowing volume, approximately half is within the Amazon river alone.

THE EARTH MOVER NUMBER

$$Em = \frac{\text{annual mass of earth moved by humans}}{\text{annual mass of earth moved by rivers}} > 15$$



The Earth Mover Number describes the mass of sediment moved by humans each year. Through construction, mining, and agriculture, humans move more than $2.5 \times 10^{14} \text{ kg / yr}$ of sediment a year (HuID: 72899, 59640, 19415, 41496). While there is uncertainty in the total mass of sediment moved through urbanization, the total mass of earth moved by humans is at least 15 times the approximate mass moved each year by the world’s rivers ($1.3 \times 10^{13} \text{ kg / yr}$ [3]). This remarkable anthropogenic action rapidly increases erosion rates, leading to increased topsoil loss and turnover, ultimately perturbing natural biogeochemical cycles.

THE ANTHROPOMASS NUMBER

$$An = \frac{\text{total anthropomass}}{\text{total biomass}} \approx 1$$



The Anthropomass Number takes stock of our material production by comparing the total quantity of human-made materials to the entirety of the biomass on planet Earth. Around 2020, total human made materials added up to the same mass as the total biomass dry weight ($\approx 1.1 \times 10^{15} \text{ kg}$ [4]). Concretes and aggregates (such as gravel) dominate the anthropomass, with bricks and asphalt coming in a distant second. Despite their ubiquity, plastics and metals constitute less than 10% of total anthropomass. Altogether, the total amounts to a dizzying $\approx 10^5 \text{ kg}$ of human made mass, or about 20 African bush elephants, per person on the planet.

THE SOLAR NUMBER

$$Su = \frac{\text{annual human power usage}}{\text{annual incident solar power}} \approx 0.0001$$



The Solar Number puts the 20 TW power consumption of human activities (HuID: 31373; 85317) in relief by comparing it to the power incident on our planet from the sun. While humans derive biological energy from food, we derive mechanical and electrical power from various fuel sources like coal, oil, natural gas, and fissile nuclear material. Current global human power usage amounts to an enormous 18 terawatts ($18 \times 10^{12} \text{ W}$) and only $\approx 1\%$ comes from solar power (0.12 – 0.20 TW, HuID: 99885; 58303). However, the incident power from the sun dwarfs this number. Of course, it is unlikely that we could ever harness 100% of solar energy incident on our planet, but capturing even 1% of this energy would provide 100 times more power than we currently use.

THE CO₂ NUMBER

$$CO_2 = \frac{\text{annual mass of anthropogenic } CO_2}{\text{annual mass of naturally removed } CO_2} \approx 2$$



The CO₂ Number compares the annual amount of human-caused CO₂ emissions to the mass of CO₂ naturally removed from the atmosphere each year. There are many climate-related consequences of increasing CO₂ emissions. Beyond accelerating climate change, $\approx 25\%$ of CO₂ released into the atmosphere is absorbed by the oceans, making them appreciably more acidic over time. In recent years, human activities, including burning fossil fuels and making concrete, have led to the release of $\approx 4 \times 10^{13} \text{ kg}$ of CO₂ (HuID: 54608, 24789) into the atmosphere each year. While many natural processes like volcanoes and wildfires release CO₂, they are generally accompanied by corresponding sinks that remove even more CO₂, like plant photosynthesis. Once all natural processes have been accounted for, a net natural sink of $\approx 2 \times 10^{13} \text{ kg}$ of CO₂ per year remains (HuID: 52670). Thus, the CO₂ number quantifies the extent to which human emissions outpace the natural removal of CO₂.

THE METHANE NUMBER

$$Me = \frac{\text{annual mass of anthropogenic } CH_4}{\text{annual mass of natural } CH_4} \approx 1$$


The CH₄ Number sheds light on the anthropogenic contribution to methane emissions. While CO₂ is the most often discussed greenhouse gas, human activities also release substantial amounts of methane (CH₄), an even more potent greenhouse gas than CO₂. Anthropogenic methane emissions result from fossil fuel extraction, ruminant livestock (mostly cows), rice cultivation, and other sources, totaling $\approx 3\text{--}4 \times 10^9 \text{ kg per year}$ (HuID: 96837). Natural emissions of CH₄, stemming mostly from wetlands and other anaerobic environments, produce a comparable amount ($\approx 2\text{--}4 \times 10^9 \text{ kg / yr}$) to anthropogenic emissions (HuID: 56405). Both of these amounts are estimates from models, due to their uncertainty, we report that anthropogenic and natural methane emissions are approximately equal in magnitude.

THE EXTINCTION NUMBER

$$Ex = \frac{\text{number of known animal extinctions}}{\text{number of expected natural animal extinctions}} > 10$$


The Extinction Number quantifies the deleterious effect that human action has on other species. Over the past 500 years, far more animal species have gone extinct than would be expected due to natural processes. Since 1500, at least 760 animal species have gone extinct (HuID: 44641). Recent estimates of ancient rates of animal extinction predict that tenfold fewer (≈ 50) species would have gone extinct over the same period in the absence of humans [5]. It’s important to emphasize that these data are incomplete and reflect only a fraction of species that have been assessed for conservation status. **The Extinction Number** therefore represents a lower bound on the degree of modern species loss, with the true value likely being much higher.

The Anthropocene by the Numbers — Supporting Information

About: Here, we present citations and notes corresponding to each quantity assessed here. Each value presented on page 1 is assigned a Human Impacts Database identifier (HuID), accessible via anthronumbers.org. When possible, primary data sources have been collated and stored as files in comma-separated-value (csv) format on the GitHub repository associated with this snapshot, accessible via DOI: 10.5281/zenodo.4453277 and https://github.com/rpgroup-pboc/human_impacts

A SURFACE TEMPERATURE

Surface temperature change relative to 1850–1900 average $\approx 1 - 1.4^\circ\text{C}$ HuID: 79598, 76539, 12147

Data Source(s): HadCRUT.4.6 (Morice et al., 2012, DOI: 10.1029/2011JD017187), GISTEMP v4 (GISTEMP Team, 2020: GISS Surface Temperature Analysis (GISTEMP), version 4. NASA Goddard Institute for Space Studies. Dataset accessed 2020-12-17 at <https://data.giss.nasa.gov/gistemp/> & Lenssen et al., 2019, DOI: 10.1029/2018JD029522) and NOAAGlobalTemp v5 (Zhang et al., 2019, DOI: 10.1029/2019EO128229) datasets. **Notes:** The global mean surface temperature captures near-surface air temperature over the planet's land and ocean surface. The value reported represents the spread of the three estimates and their 95% confidence intervals. Temperature changes from all three datasets are expressed relative to the 1850–1900 average temperature from the HadCRUT.4.6 dataset. Since data for the period 1850–1880 are missing in GISTEMP v4 and NOAAGlobalTemp v5, data are centered by setting the 1880–1900 mean of all datasets to the HadCRUT.4.6 mean over the same period.

B ANNUAL ICE MELT

glaciers $= (3.0 \pm 1.2) \times 10^{11} \text{ m}^3 / \text{yr}$ HuID: 32459

Data Source(s): Intergovernmental Panel on Climate Change (IPCC) 2019 Special Report on the Ocean and Cryosphere in a Changing Climate. Table 2.A.1 on pp. 199–202. **Notes:** Value corresponds to the trend of annual glacial ice volume loss (reported as ice mass loss) from major glacierized regions (2006–2015) based on aggregation of observation methods (original data source: Zemp et al. 2019, DOI:10.1038/s41586-019-1071-0) with satellite gravimetric observations (original data source: Wouters et al. 2019, DOI:10.3389/feart.2019.00096). Ice volume loss was calculated from ice mass loss assuming a standard pure ice density of $920 \text{ kg} / \text{m}^3$. Uncertainty represents a 95% confidence interval calculated from standard error propagation of the 95% confidence intervals reported in the original sources assuming them to be independent.

ice sheets $= (4.7 \pm 0.4) \times 10^{11} \text{ m}^3 / \text{yr}$ HuID: 95798, 93137

Data Source(s): D. N. Wiese et al. 2019 JPL GRACE and GRACE-FO Mascon Ocean, Ice, and Hydrology Equivalent HDR Water Height RLO6M CRI Filtered Version 2.0, Ver. 2.0, PO.DAAC, CA, USA. Dataset accessed [2020-Aug-10]. DOI: 10.5067/TEM-SC-3JM62

Notes: Value corresponds to the trends of combined annual ice volume loss (reported as ice mass loss) from the Greenland and Antarctic Ice Sheets (2002–2020) measured by satellite gravimetry. Ice volume loss was calculated from ice mass loss assuming a standard pure ice density of $920 \text{ kg} / \text{m}^3$. Uncertainty represents one standard deviation and considers only propagation of monthly uncertainties in measurement.

Arctic sea ice $= (3.0 \pm 1.0) \times 10^{11} \text{ m}^3 / \text{yr}$ HuID: 89520

Data Source(s): PIOMAS Arctic Sea Ice Volume Reanalysis, Figure 1 of webpage as of October 31, 2020. Original method source: Schweiger et al. 2011, DOI:10.1029/2011JC007084

Notes: Value reported corresponds to the trend of annual volume loss from Arctic sea ice (1979–2020). The uncertainty in the trend represents the range in trends calculated from three ice volume determination methods.

C SEA ICE EXTENT

extent loss at yearly maximum cover (September) $\approx 8.4 \times 10^{10} \text{ m}^2 / \text{yr}$ HuID: 33993

extent loss at yearly minimum cover (March) $\approx 4.0 \times 10^{10} \text{ m}^2 / \text{yr}$ HuID: 87741

average annual extent loss $= 5.5 \pm 0.2 \times 10^{10} \text{ m}^2 / \text{yr}$ HuID: 70818

Data Source(s): Comiso et al. 2017, DOI:10.1002/2017JC012768. Fetterer et al. 2017, updated daily. Sea Ice Index, Version 3, Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center, DOI:10.7265/N5K072F8, [Accessed 2020-Oct-19]. **Notes:** Sea ice extent refers to the area of the sea with $> 15\%$ ice coverage. Annual value corresponds to the linear trend of annually averaged Arctic sea ice extent from 1979–2015 (Comiso et al. 2017) calculated from four different methods. This is in good agreement with the linear trend of annual extent loss calculated by averaging over every month in a given year ($5.5 \times 10^{10} \text{ m}^2 / \text{yr}$ HuID: 66277). The minimum cover extent loss corresponds to the linear trend of Arctic sea ice extent in September from 1979–2020 and the maximum cover extent loss corresponds to the linear trend of sea ice extent in March from 1979–2020. The Antarctic sea ice extent trend is not shown because a significant long-term trend over the satellite observation period is not observed and short-term trends are not yet identifiable.

D MATERIAL PRODUCTION

concrete production $\approx (2 - 3) \times 10^{13} \text{ kg} / \text{yr}$ HuID: 25488, 81346, 16995

Data Source(s): United States Geological Survey (USGS), Mineral Commodity Summaries 2020, pp. 42–43, DOI:10.3133/mcs2020. Miller et al. 2016, Table 1, DOI:10.1088/1748-9326/11/7/074029. Monteiro et al. 2017, DOI:10.1038/nmat4930. Krausmann et al. 2017, DOI:10.1073/pnas.1613773114 **Notes:** Concrete is formed when aggregate material is bonded together by hydrated cement. The USGS reports the mass of cement produced in 2019 as $4.1 \times 10^{12} \text{ kg}$ in 2019. As most cement is used to form concrete, cement production can be used to estimate concrete mass using a multiplicative conversion factor of 7 (Monteiro et al.). Miller et al. report that the cement, aggregate and water used in concrete in 2012 sum to $2.3 \times 10^{13} \text{ kg}$. Krausmann et al. report an estimated value from 2010 based on a material input, stocks, and outputs model. The value is net annual addition to concrete stocks plus annual waste and recycling to estimate gross production of concrete.

steel production $= (1.4 - 1.9) \times 10^{13} \text{ kg} / \text{yr}$ HuID: 51453, 44894, 85981

Data Source(s): United States Geological Survey (USGS), Mineral Commodity Summaries 2020, pp. 82–83, DOI:10.3133/mcs2020. World Steel Association, World Steel in Figures 2020, p. 6. Krausmann et al. 2017, DOI:10.1073/pnas.1613773114 **Notes:** Crude steel includes stainless steels, carbon steels, and other alloys. The USGS reports the mass of crude steel produced in 2019 as 1900 megatonnes (Mt). The World Steel Association reports a production value of 1869 Mt in 2019. Krausmann et al. report an estimated value from 2010 based on a material input, stocks, and outputs model. The value is net annual addition to steel stocks plus annual waste and recycling to estimate gross production of steel.

plastic production $\approx 4 \times 10^{11} \text{ kg} / \text{yr}$ HuID: 97241, 25437

Data Source(s): Geyer et al. 2017, Table S1, DOI:10.1126/sciadv.1700782; Krausmann et al. 2017, DOI:10.1073/pnas.1613773114. **Notes:** Value represents the approximate sum total global production of plastic fibers and plastic resin during the calendar year of 2015. Comprehensive data about global plastic production is sorely lacking. Geyer et al. draw data from various industry groups to estimate total production of different polymers and additives. Some of the underlying data is not publicly available, and data from financially-interested parties is inherently suspect. Krausmann et al. report an estimated value from 2010 based on a material input, stocks, and outputs model. The value is net annual addition to stocks plus annual waste and end-of-life recycling to estimate gross production of plastics.

E LIVESTOCK POPULATION

chicken standing population $\approx 2.5 \times 10^{10}$ HuID: 94934

cattle standing population $\approx 1.5 \times 10^9$ HuID: 92006

swine standing population $\approx 1 \times 10^9$ HuID: 21368

all livestock standing population $\approx 3 \times 10^{10}$ HuID: 43599

Data Source(s): Food and Agriculture Organization (FAO) of the United Nations Statistical Database (2020) — Live Animals. **Notes:** Counts correspond to the estimated standing populations in 2018. Values are reported directly by countries. The FAO uses non-governmental statistical sources to address uncertainty and missing (non-reported) data. Reported values are therefore approximations.

F SYNTHETIC NITROGEN FIXATION

annual mass of synthetically fixed nitrogen $\approx 1.5 \times 10^{11} \text{ kg N} / \text{yr}$ HuID: 60580, 61614

Data Source(s): United States Geological Survey (USGS), Mineral Commodity Summaries 2020, pp. 116–117, DOI:10.3133/mcs2020. International Fertilizer Association (IFA) Statistical Database (2020) — Ammonia Production & Trade Tables by Region. Smith et al. 2020, DOI: 10.1039/c9ee02873k. **Notes:** Ammonia (NH_3) produced globally is compiled by the USGS and IFA from major factories that report output. The USGS estimates the approximate mass of nitrogen in ammonia produced in 2019 as $1.5 \times 10^{11} \text{ kg N}$ and the International Fertilizer Association reports a production value of $1.5 \times 10^{11} \text{ kg N}$ in 2019. Nearly all of this mass is produced by the Haber-Bosch process ($>96\%$, Smith et al. 2020). In the United States most of this mass is used for fertilizer, with the remainder being used to synthesize nitrogen-containing chemicals including explosives, plastics, and pharmaceuticals ($\approx 88\%$, USGS Mineral Commodity Summaries 2020).

G OCEAN ACIDITY

surface ocean $[\text{H}^+] \approx 0.2$ parts per billion HuID: 90472

annual change in $[\text{H}^+] = 0.36 \pm 0.03\%$ HuID: 19394

Data Source(s): Figures 1–2 of European Environment Agency report CLIM 043 (2020). Original data source of the report is “Global Mean Sea Water pH” from Copernicus Marine Environment Monitoring Service. **Notes:** Reported value is calculated from the global average annual change in pH over years 1985–2018. The average oceanic pH was ≈ 8.057 in 2018 and decreases annually by ≈ 0.002 units, giving a change in $[\text{H}^+]$ of roughly $10^{-8.056} - 10^{-8.057} \approx 4 \times 10^{-11} \text{ mol/L}$ or about 0.4% of the global average. $[\text{H}^+]$ is calculated as $10^{-\text{pH}} \approx 10^{-8} \text{ mol/L}$ or 0.2 parts per billion (ppb) which is calculated by noting that $[\text{H}_2\text{O}] \approx 55 \text{ mol} / \text{L}$. Uncertainty for annual change is the standard error of the mean.

H LAND USE

agricultural $\approx 5 \times 10^{13} \text{ m}^2$ HuID: 29582

Data Source(s): Food and Agriculture Organization (FAO) of the United Nations Statistical Database (2020) — Land Use. **Notes:** Agricultural land is defined as all land that is under agricultural management including pastures, meadows, permanent crops, temporary crops, land under fallow, and land under agricultural structures (such as barns). Reported value corresponds to 2017 estimates by the FAO.

urban $\approx (6 - 8) \times 10^{11} \text{ m}^2$ HuID: 41339, 39341

Data Source(s): Florczyk et al. 2019 (<https://tinyurl.com/yxxxtgll>) and Table 3 of Liu et al. 2018 DOI: 10.1016/j.rse.2018.02.055. **Notes:** Urban land area is determined from satellite imagery. An area is determined to be “urban” if the total population is greater than 5,000 and has a minimum population density of 300 people per km^2 . Reported value gives the range of recent measurements of $\approx 6.5 \times 10^{11} \text{ m}^2$ (2015) and $\approx (7.5 \pm 1.5) \times 10^{11} \text{ m}^2$ (2010) from Florczyk et al. 2019 and Liu et al. 2018, respectively.

I RIVER FRAGMENTATION

global fragmented river volume $\approx 6 \times 10^{11} \text{ m}^3$ HuID: 61661

Data Source(s): Grill et al. 2019 DOI: 10.1038/s41586-019-1111-9. **Notes:** Value corresponds to the water volume contained in rivers that fall below the connectivity threshold required to classify them as free-flowing. Value considers only rivers with upstream catchment areas greater than 10 km^2 or discharge volumes greater than 0.1 m^3 per second. The ratio of global river volume in disrupted rivers to free-flowing rivers is approximately 0.9. The exact value depends on the cutoff used to define a “free-flowing” river. We direct the reader to the source for thorough detail.

J HUMAN POPULATION

urban-dwelling fraction of population $\approx 55\%$ HuID: 93995

total population $\approx 7.6 \times 10^9$ HuID: 85255

Data Source(s): Food and Agricultural Organization (FAO) of the United Nations Report on Annual Population, 2019. **Notes:** Value for total population in 2018 comes from a combination of direct population reports from country governments as well as inferences of underreported or missing data. The definition of “urban” differs between countries and the data does not distinguish between urban and suburban populations despite substantive differences between these land uses (Jones and Kammen 2013, doi: 10.1021/es4034364). As explained by the United Nations population division, “When the definition used in the latest census was not the same as in previous censuses, the data were adjusted whenever possible so as to maintain consistency.” Rural population is computed from this fraction along with the total human population, implying that the total population is composed only of “urban” and “rural” communities.

K GREENHOUSE GAS EMISSIONS

anthropogenic $\text{CO}_2 = (4.25 \pm 0.33) \times 10^{13} \text{ kg CO}_2 / \text{yr}$ HuID: 24789, 54608, 98043

Data Source(s): Table 6 of Friedlingstein et al. 2019, DOI: 10.5194/essd-11-1783-2019. Original data sources relevant to this study compiled in Friedlingstein et al.: 1) Giffillan et al. <https://energy.appstate.edu/CDIAC> 2) Average of two bookkeeping models: Houghton and Nassikas 2017 DOI: 10.1002/2016GB005546; Hansis et al. 2015 DOI: 10.1002/2014GB004997) Dlugokencky and Tans, NOAA/GML <https://www.esrl.noaa.gov/gmd/ccgg/trends/>. **Notes:** Value corresponds to total CO_2 emissions from fossil fuel combustion, industry (predominantly cement production), and land-use change during calendar year 2018. Emissions from land-use change are due to the burning or degradation of plant biomass. In 2018, $1.88 \times 10^{13} \text{ kg CO}_2 / \text{yr}$ accumulated in the atmosphere, reflecting the balance of emissions and CO_2 uptake by plants and oceans. Uncertainty corresponds to one standard deviation.

The Anthropocene by the Numbers — Supporting Information

K GREENHOUSE GAS EMISSIONS (CONTINUED)

anthropogenic $\text{CH}_4 = (3.4 - 3.9) \times 10^{11} \text{ kg CH}_4 / \text{yr}$ [HuID: 96837, 30725](#)
Data Source(s): Table 3 of Saunio, et al. 2020. DOI: 10.5194/essd-12-1561-2020. **Notes:** Value corresponds to 2008–2017 decadal average mass of CH_4 emissions from anthropogenic sources. Includes emissions from agriculture and landfill, fossil fuels, and burning of biomass and biofuels, but other inventories of anthropogenic methane emissions are also considered. Reported range represents the minimum and maximum estimated emissions from a combination of “bottom-up” and “top-down” models.

anthropogenic $\text{N}_2\text{O} = 1.1 (+0.6, -0.5) \times 10^{10} \text{ kg N}_2\text{O} / \text{yr}$ [HuID: 44575](#)

Data Source(s): Table 1 of Tian, H., et al. 2020. DOI: 10.1038/s41586-020-2780-0. **Notes:** Value corresponds to annualized N_2O emissions from anthropogenic sources in the years 2007–2016. The value reported in the source is 7.3 (4.2, 11.4) Tg N / year. This is converted to a mass of N_2O using the fact that $\text{N} \approx 14/22$ of the mass of N_2O . Reported value is mean with the uncertainty bounds (+, -) representing the maximum and minimum values observed in the 2007–2016 time period.

L WATER WITHDRAWAL

agricultural withdrawal = $1.3 \times 10^{12} \text{ m}^3 / \text{yr}$ [HuID: 84545, 43593, 95345](#)

industrial withdrawal = $5.9 \times 10^{11} \text{ m}^3 / \text{yr}$ [HuID: 27142](#)

domestic withdrawal = $5.4 \times 10^{10} \text{ m}^3 / \text{yr}$ [HuID: 69424](#)

total withdrawal = $(1.7 - 2.2) \times 10^{12} \text{ m}^3 / \text{yr}$ [HuID: 27342, 68004](#)

Data Source(s): Figure 1 of Qin et al. 2019. DOI: 10.1038/s41893-019-0294-2. AQUASTAT Main Database, Food and Agriculture Organization of the United Nations **Notes:** “Agricultural” and “total” withdrawal include one value from Qin et al. (who reports “consumption”) and one value from the AQUASTAT database. Industrial water withdrawal is from AQUASTAT and domestic withdrawal value is from Qin et al. Values in AQUASTAT are self-reported by countries and have missing values from some countries, probably accounting for a few percent underreporting. All values represent withdrawals. For agricultural and domestic, water withdrawal is assumed to be the same as water consumption as reported in Qin et al.

M SEA LEVEL RISE

added water = $1.97 (+0.36, -0.34) \text{ mm} / \text{yr}$ [HuID: 97108](#)

thermal expansion = $1.19 (+0.25, -0.24) \text{ mm} / \text{yr}$ [HuID: 97688](#)

total observed sea-level rise = $3.35 (+0.47, -0.44) \text{ mm} / \text{yr}$ [HuID: 81373](#)

Data Source(s): Table 1 of Frederikse et al. 2020. DOI: 10.1038/s41586-020-2591-3. **Notes:** Values correspond to the average global sea level rise for the years 1993 – 2018. “Added water” (barystatic) change includes effects from meltwater from glaciers and ice sheets, added mass from sea-ice discharge, and changes in the amount of terrestrial water storage. Thermal expansion accounts for the volume change of water with increasing temperature. Values for “thermal expansion” and “added water” come from direct observations of ocean temperature and gravimetry/altimetry, respectively. Total sea level rise is the observed value using a combination of measurement methods. “Other sources” reported on page 1 accounts for observed residual sea level rise not attributed to a source in the model. Values in brackets correspond to the upper and lower bounds of the 90% confidence interval.

N TOTAL POWER USE

global power use $\approx 19 - 20 \text{ TW}$ [HuID: 31373, 85317](#)

Data Source(s): bp Statistical Review of World Energy, 2020; U.S. Energy Information Administration, 2020. **Notes:** Value represents the sum of total primary energy consumed from oil, natural gas, coal, and nuclear energy and electricity generated by hydroelectric and other renewables. Value is calculated using annual primary energy consumption as reported in data sources assuming uniform use throughout a year, yielding $\approx 19 - 20 \text{ TW}$.

O TREE COVERAGE AREA LOSS

commodity-driven deforestation = $(5.7 \pm 1.1) \times 10^{10} \text{ m}^2 / \text{yr}$ [HuID: 96098](#)

forestry = $(5.4 \pm 0.8) \times 10^{10} \text{ m}^2 / \text{yr}$ [HuID: 38352](#)

urbanization = $(2 \pm 1) \times 10^9 \text{ m}^2 / \text{yr}$ [HuID: 19429](#)

shifting agriculture = $(7.5 \pm 0.9) \times 10^{10} \text{ m}^2 / \text{yr}$ [HuID: 24388](#)

wildfire = $(7.2 \pm 1.3) \times 10^{10} \text{ m}^2 / \text{yr}$ [HuID: 92221](#)

total loss $\approx 2 \times 10^{11} \text{ m}^2 / \text{yr}$ [HuID: 78576](#)

Data Source(s): Table 1 of Curtis et al. 2018 DOI: 10.1126/science.aau3445. Hansen et al. 2013 DOI: 10.1126/science.1244693. Global Forest Watch, 2020. Reported values in source correspond to total loss from 2001 – 2015. Values given are averages over this 15 year window. **Notes:** Commodity-driven deforestation is “long-term, permanent, conversion of forest and shrubland to a non-forest land use such as agriculture, mining, or energy infrastructure.” Forestry is defined as large-scale operations occurring within managed forests and tree plantations with evidence of forest regrowth in subsequent years. Urbanization converts forest and shrubland for the expansion and intensification of existing urban centers. Disruption due to “shifting agriculture” is defined as “small- to medium-scale forest and shrubland conversion for agriculture that is later abandoned and followed by subsequent forest regrowth”. Disruption due to wildfire is “large-scale forest loss resulting from the burning of forest vegetation with no visible human conversion or agricultural activity afterward”. Uncertainty corresponds to the 95% confidence interval. Uncertainty is approximate for “urbanization” as the source reports an ambiguous error of “ $\pm <1\%$ ”.

P POWER FROM FOSSIL FUELS

natural gas = $4.5 - 4.8 \text{ TW}$ [HuID: 49947, 86175](#)

oil = $6.1 - 6.6 \text{ TW}$ [HuID: 4121, 39756](#)

coal = $5.0 - 5.5 \text{ TW}$ [HuID: 10400, 60490](#)

total = $16 - 17 \text{ TW}$ [HuID: 29470, 29109](#)

Data Source(s): bp Statistical Review of World Energy, 2020. U.S. Energy Information Administration, 2020. **Notes:** Values are self-reported by countries. Values from bp Statistical Review correspond to 2019 whereas values from the EIA correspond to 2018 estimates. Reported TW are computed from primary energy (e.g. kg coal) units assuming uniform use throughout the year. Oil volume includes crude oil, shale oil, oil sands, condensates, and natural gas liquids separate from specific natural gas mining. Natural gas value excludes gas flared or recycled and includes natural gas produced for gas-to-liquids transformation. Coal value includes 2019 value exclusively for solid commercial fuels such as bituminous coal and anthracite, lignite and subbituminous coal, and other solid fuels. This includes coal used directly in power production as well as coal used in coal-to-liquids and coal-to-gas transformations.

Q POWER FROM RENEWABLE RESOURCES

wind $\approx 0.36 - 0.39 \text{ TW}$ [HuID: 30581, 85919](#)

solar $\approx 0.18 - 0.20 \text{ TW}$ [HuID: 99885, 58303](#)

hydroelectric = 1.2 TW [HuID: 15765, 50558](#)

total renewable power $\approx 1.9 - 2.1 \text{ TW}$ [HuID: 75741, 20246](#)

Data Source(s): bp Statistical Review of World Energy, 2020. U.S. Energy Information Administration, 2020. **Notes:** Reported values correspond to estimates for the 2019 and 2018 calendar years for bp and EIA sources, respectively. Renewable resources are defined as wind, geothermal, solar, biomass and waste. Hydroelectric, while presented here, is not defined as a renewable in the bp dataset. All values are reported as input-equivalent energy, meaning the input energy that would have been required if the power was produced by fossil fuels. BP reports that fossil fuel efficiency used to make this conversion was $\approx 40\%$ in 2017.

R FOSSIL FUEL EXTRACTION

volume of natural gas = $(3.9 - 4.0) \times 10^{12} \text{ m}^3 / \text{yr}$ [HuID: 11468, 20532](#)

volume of oil = $(5.5 \pm 5.8) \times 10^9 \text{ m}^3 / \text{yr}$ [HuID: 66789, 97719](#)

mass of coal = $(7.8 - 8.1) \times 10^{12} \text{ kg} / \text{yr}$ [HuID: 78435, 48928](#)

Data Source(s): bp Statistical Review of World Energy, 2020. U.S. Energy Information Administration, 2020. **Notes:** Oil volume includes crude oil, shale oil, oil sands, condensates, and natural gas liquids separate from specific natural gas mining. Natural gas value excludes gas flared or recycled and includes natural gas produced for gas-to-liquids transformation. Coal value includes solid commercial fuels such as bituminous coal, anthracite, lignite, subbituminous coal, and other solid fuels. All values from bp Statistical Review correspond to 2019 whereas values from the EIA correspond to 2018 estimates.

S OCEAN WARMING

heat uptake by ocean $\approx 346 \pm 51 \text{ TW}$ [HuID: 94108](#)

upper ocean (0 – 700 m) temperature increase since 1960 = $0.18 - 0.2 \text{ }^\circ\text{C}$ [HuID: 69674, 72086](#)

Data Source(s): Table S1 of Cheng et al. 2017. doi: 10.1126/sciadv.1601545. NOAA National Centers for Environmental Information, 2020. doi: 10.1029/2012GL051106. **Notes:** Heat uptake reported is the average over time period 1992–2015 with 95% confidence intervals. Range of temperatures reported captures the 95% confidence interval of temperature increase for the period 2015–2019 with respect to the 1958–1962 mean. Temperature change is considered in the upper 700 m because sea surface temperatures have high decadal variability and are a poor indicator of ocean warming; see Roemmich et al. 2015, doi: 10.1038/NCLIMATE2513.

T POWER FROM NUCLEAR FISSION

nuclear power $\approx 0.79 - 0.89 \text{ TW}$ [HuID: 48387](#)

Data Source(s): bp Statistical Review of World Energy, 2020. U.S. Energy Information Administration, 2020. **Notes:** Values are self-reported by countries and correspond to estimates for 2019 and 2018 calendar year for bp and EIA data, respectively. Values are reported as “input-equivalent” energy, meaning the energy needed to produce a given amount of power if the input were a fossil fuel, which is converted to TW here. This is calculated by multiplying the given power by a conversion factor representing the efficiency of power production by fossil fuels. In 2017, this factor was about 40%.

U NUCLEAR FALLOUT

anthropogenic ^{239}Pu and ^{240}Pu from weapons testing $\approx 1.4 \times 10^{11} \text{ kg} / \text{yr}$ [HuID: 42526](#)

Data Source(s): Table 1 in Hancock et al. 2014 doi: 10.1144/SP395.15. Fallout in activity from UNSCEAR 2000 Report on Sources and Effects of Ionizing Radiation Report to the UN General Assembly -- Volume 1. **Notes:** The approximate mass of Plutonium isotopes ^{239}Pu and ^{240}Pu released into the atmosphere from the ≈ 500 above-ground nuclear weapons tests conducted between 1945 and 1980. Naturally occurring ^{239}Pu and ^{240}Pu are rare, meaning that nearly all contemporary labile plutonium comes from human production. (Taylor 2001, doi: 10.1016/S1569-4860(01)80003-6) The total mass of radionuclides released is $\approx 3300 \text{ kg}$ with a combined radioactive fallout of $\approx 11 \text{ PBq}$. These values do not represent the entire $^{239+240}\text{Pu}$ globally distributed mass as it excludes non-weapons sources.

V CONTEMPORARY EXTINCTION

animal species extinct since 1500 > 750 [HuID: 44641](#)

plant species extinct since 1500 > 120 [HuID: 86866](#)

Data Source(s): The IUCN Red List of Threatened Species. Version 2020–2. **Notes:** Values correspond to absolute lower-bound count of animal extinctions caused over the past ≈ 520 years. Of the predicted ≈ 8 million animal species, the IUCN databases catalogues only $\approx 900,000$ with only $\approx 75,000$ being assigned a conservation status. Representation of plants and fungi is even more sparse with only $\approx 40,000$ and ≈ 285 being assigned a conservation status, respectively. The number of extinct animal species is undoubtedly higher than these reported values, as signified by an inequality symbol ($>$).

W EARTH MOVING

waste and overburden from coal mining $\approx 6.5 \times 10^{13} \text{ kg} / \text{yr}$ [HuID: 72899](#)

earth moved from urbanization $> 1.4 \times 10^{14} \text{ kg} / \text{yr}$ [HuID: 59640](#)

Data Source(s): Supplementary table 1 of Cooper et al. 2018. DOI: doi.org/gfwfhnd. **Notes:** Coal mining waste and overburden mass is calculated given commodity-level stripping ratios (mass of overburden/waste per mass of coal resource mined) and reported values of global coal production by type. Urbanization mass is presented as a lower bound estimate of the mass of earth moved from global construction projects. This comes from a conservative estimate that the ratio of the mass of earth moved per mass of cement/concrete used in construction globally is 2:1. This value is highly context dependent and we encourage the reader to read the source material for a more thorough description of this estimation.

erosion from agricultural land $> 1.2 - 2.4 \times 10^{13} \text{ kg} / \text{yr}$ [HuID: 19415, 41496](#)

Data Source(s): Pg. 377 of Wang and Van Oost 2019. DOI: 10.1177/0959683618816499. Pg. 21996 of Borrelli et al. 2020 DOI: 10.1073/pnas.2001403117. **Notes:** Cumulative sediment mass loss over history of human agriculture due to accelerated erosion is estimated to be $\approx 30,000 \text{ Gt}$. Recent years have an estimated erosion rate ranging from $12 \text{ Pg} / \text{yr}$ (Wang and Van Oost) to $\approx 24 \text{ Pg} / \text{yr}$ (Borrelli et al.). Values come from computational models conditioned on time-resolved measurements of sediment deposition in catchment basins.